

Spectral Theory on Commutative Kreĭn C*-algebras

Pichkitti Bannangkoon^{a*}, Paolo Bertozzini^b, Wicharn Lewkeeratiyutkul^a

^a*Department of Mathematics and Computer Science, Faculty of Science,
Chulalongkorn University, Bangkok 10330, Thailand*

e-mails: pb_math@yahoo.com Wicharn.L@chula.ac.th

^b*Department of Mathematics and Statistics, Faculty of Science and Technology,
Thammasat University, Bangkok 12121, Thailand*

e-mail: paolo.th@gmail.com

08 December 2013.[†]

Abstract

A Banach involutive algebra $(A, *)$ is called a Kreĭn C*-algebra if there is a fundamental symmetry of $(A, *)$, i.e., $\alpha \in \text{Aut}(A, *)$ such that $\alpha^2 = 1_A$ and $\|\alpha(x^*)x\| = \|x\|^2$ for all $x \in A$. Using α , we can decompose $A = A_+ \oplus A_-$ where $A_+ = \{x \in A \mid \alpha(x) = x\}$ and $A_- = \{x \in A \mid \alpha(x) = -x\}$. The even part A_+ is a C*-algebra and the odd part A_- is a Hilbert C*-bimodule over the even part A_+ . The ultimate goal is to develop a spectral theory for commutative unital Kreĭn C*-algebras when the odd part is a symmetric imprimitivity bimodule over the even part and there exists a suitable “exchange symmetry” ϵ between A_+ and A_- .

Keywords: Spectral Theory, Kreĭn C*-algebras.

MSC-2010: 47B50, 46C20.

1 Introduction

The study of vector spaces equipped with “inner products” that are non-necessarily positive-definite has always been a theme of extreme importance in relativistic physics starting probably with the work of H.Minkowski [M]. Complex vector spaces with indefinite sesquilinear forms have been introduced in relativistic quantum field theory by P.Dirac [D], W.Pauli [P] and then used by S.Gupta [Gu] K.Bleuler [B], although their mathematical definition has been given later by L.Pontrjagin [Po]. Complete “indefinite” inner product spaces, called Kreĭn spaces, have been introduced by Ju.Ginzburg [G] and E.Scheibe [S] and the study of their properties has been undertaken by several Russian mathematicians.

Although algebras of continuous operators on Kreĭn spaces have been around for some time, a first definition of abstract Kreĭn C*-algebra has been provided only recently by K.Kawamura [K1, K2]. Kreĭn C*-algebras are somehow expected to play some role in a “semi-Riemannian” version of A.Connes non-commutative geometry [C] (see A.Strohmaier [St], M.Pasche-A.Rennie-R.Verch [PRV] for details) and for this reason it should be of some interest to develop a spectral theory that is suitable for them.

It is the purpose of this paper to introduce a simple spectral theory for the special class of Kreĭn C*-algebras that decompose, via a fundamental symmetry, in direct sum $A = A_+ \oplus A_-$ with A_+

^{*}Current Address: Department of Mathematics, Penn State University, University Park, State College, PA 16802, USA.

[†]This is a reformatted version for arXiv of an original paper presented at the “14th Annual Thai Mathematical Meeting” in Prince of Songkla University Pattani Campus on 5-6 March 2009 and appeared in: 14th Annual Meeting in Mathematics 2009, Collection of Full Papers 38-55.

commutative C^* -algebra and A_- symmetric imprimitivity (anti-)Hilbert C^* -bimodule over A_+ and that are equipped with an odd symmetry exchanging A_+ and A_- . Our main result (Theorem 3.27) is that every such Kreĭn C^* -algebra turns out to be isomorphic (via Gel'fand transform) to an algebra of continuous functions with values in a very elementary Kreĭn C^* -algebra $\mathbb{K}$ defined in detail in Theorem 3.10.

The main result presented here can actually be also obtained in at least a few other ways that we briefly describe here below:

- For a symmetric imprimitivity commutative unital Kreĭn C^* -algebra A , the even part A_+ , as a commutative unital C^* -algebra, is isomorphic to the algebra of sections of a trivial complex line bundle over the Gel'fand spectrum $\Omega(A_+)$. The odd part A_- is a symmetric imprimitivity Hilbert C^* -bimodule over A_+ and, making use of the spectral theorem for imprimitivity Hilbert C^* -bimodules developed in [BCL2], it is isomorphic to the bimodule of sections of a complex line bundle over the same $\Omega(A_+)$. Under the existence of an odd symmetry on A , the Whitney sum of the previous two line bundles turns out to be a bundle of rank-one Kreĭn C^* -algebras isomorphic to $\mathbb{K}$ over $\Omega(A_+)$.
- Although we are not aware now of a specific reference to a Gel'fand theorem, once a specific fundamental symmetry/odd symmetry α, ϵ has been chosen on A , the Kreĭn C^* -algebra becomes completely equivalent to a $\mathbb{Z}_2$ -graded C^* -algebra with commutative even part and for such elementary C^* -algebras spectral results are for sure obtainable as special cases from the general theory of C^* -dynamical systems.
- To every unital Kreĭn C^* -algebra equipped with a given fundamental symmetry, we can always associate a C^* -category with two objects. In the case of imprimitivity commutative unital Kreĭn C^* -algebras, such C^* -category will be commutative and full according to the definition provided in [BCL1] and our spectral theorem can be recovered as a trivial application of the general spectral theory for commutative full C^* -categories developed in [BCL1].

The techniques utilized here in the proof of this result are essentially an adaptation of those developed in [BCL1] for a spectral theory of commutative full C^* -categories. Our choice to develop a completely independent proof of the result can be justified from the desire to test and calibrate some of the general techniques introduced in [BCL1] in a simple situation that in the near future might be used as a “laboratory” for non-commutative extensions of the spectral theorem. It is expected that more powerful spectral theories for wider classes of Kreĭn C^* -algebras might be developed using a Kreĭn version of the spaceoid Fell bundle introduced in [BCL1], we hope to address this more general problem in other preprints.

2 Preliminaries

For the convenience and usefulness of the reader, we provide here some background material and beneficial theorems on the theory of C^* -algebras and Hilbert C^* -modules (see [Mu] [BCL2] for all the details).

Definition 2.1. An **algebra** over the complex numbers is a complex vector space A equipped with a binary operation (called product) $\cdot : A \times A \rightarrow A$, $\cdot : (x, y) \mapsto x \cdot y$, that is bilinear.

The algebra is called **associative** if: $x \cdot (y \cdot z) = (x \cdot y) \cdot z$.

The algebra is called **unital** if: $\exists 1_A \in A$, such that $\forall x \in A$, $x \cdot 1_A = 1_A \cdot x = x$.

Definition 2.2. An algebra is called **involutive** or also **$*$ -algebra** if it is equipped with a function $*$: $A \rightarrow A$, such that:

$$\begin{aligned} (x^*)^* &= x, \quad \forall x \in A, \\ (x \cdot y)^* &= y^* \cdot x^*, \quad \forall x, y \in A, \\ (\alpha x + \beta y)^* &= \bar{\alpha} x^* + \bar{\beta} y^*, \quad \forall \alpha, \beta \in \mathbb{C}, \forall x, y \in A. \end{aligned}$$

Definition 2.3. A **normed algebra** is an algebra A that is also a normed space and that satisfies the property: $\|x \cdot y\| \leq \|x\| \cdot \|y\|$, $\forall x, y \in A$. A **Banach algebra** is a normed algebra that, as a normed space, is complete.

Definition 2.4. A **pre-C*-algebra** is an involutive algebra A that is also a normed algebra that satisfies the property $\|x^*x\| = \|x\|^2$, $\forall x \in A$. A **C*-algebra** is a pre-C*-algebra that is also Banach algebra.

Example 2.5. The followings are examples of C*-algebras:

1. Let X be a compact Hausdorff space. Then the space $C(X)$ of all continuous complex-valued functions on X is a unital commutative C*-algebra with the following operations and norm:
 $\forall x \in X$,

$$\begin{aligned} (f + g)(x) &= f(x) + g(x), & (cf)(x) &= cf(x), \\ (fg)(x) &= f(x)g(x), & (f^*)(x) &= \overline{f(x)}, \\ \|f\| &= \sup\{|f(x)| : x \in X\}. \end{aligned}$$

2. The space $B(H)$ of bounded linear maps on a Hilbert space H is a unital C*-algebra that is non-commutative if $\dim(H) > 1$, with the following operations and norm:

$$\begin{aligned} (T + S)(x) &= T(x) + S(x), & (cT)(x) &= cT(x), \\ (T \circ S)(x) &= T(S(x)), & \|T\| &= \sup_{x \neq 0} \frac{\|T(x)\|}{\|x\|}, \end{aligned}$$

the involution T^* of T is the adjoint of T : $\langle Tx | y \rangle = \langle x | T^*y \rangle$, $\forall x, y \in H$. ┘

Definition 2.6. A **homomorphism** from an algebra A to an algebra B is a linear map $\varphi : A \rightarrow B$ such that

$$\varphi(ab) = \varphi(a)\varphi(b) \quad \forall a, b \in A.$$

A ***-homomorphism** $\varphi : A \rightarrow B$ of *-algebras A and B is a homomorphism of algebras such that $\varphi(a^*) = \varphi(a)^*$ $\forall a \in A$. If in addition φ is a bijection, it is a ***-isomorphism**.

Definition 2.7. A **character** on a commutative algebra A is a nonzero homomorphism $\tau : A \rightarrow \mathbb{C}$. We denote by $\Omega(A)$ the set of characters on A .

For each $a \in A$, define $\hat{a} : \Omega(A) \rightarrow \mathbb{C}$ by $\hat{a}(\tau) = \tau(a)$ for all $\tau \in \Omega(A)$. Equip $\Omega(A)$ with the smallest topology on $\Omega(A)$ which makes each $\hat{a}$ continuous.

Theorem 2.8. If A is a unital commutative Banach algebra, then $\Omega(A)$ is a compact Hausdorff space with respect to the topology defined above.

Theorem 2.9 (Gel'fand-Mazur). If A is a unital Banach algebra in which every non-zero element is invertible, then $A = \mathbb{C}1$.

Theorem 2.10 (Spectral theorem). If A is a non-zero unital commutative C*-algebra, then the Gelfand transform

$$\varphi : A \rightarrow C(\Omega(A)), \quad a \mapsto \hat{a}, \quad (\hat{a} : \Omega(A) \rightarrow \mathbb{C}, \quad \tau \mapsto \tau(a))$$

is an isometric *-isomorphism.

Definition 2.11. A **right R -module** E over a ring R is an Abelian group $(E, +)$ equipped with an operation $\cdot : E \times R \rightarrow E$, $\cdot : (x, a) \mapsto x \cdot a$, of right multiplication by elements of the ring R , that satisfies the following properties:

$$\begin{aligned} x \cdot (a + b) &= (x \cdot a) + (x \cdot b), \quad \forall x \in E, \forall a, b \in R, \\ (x + y) \cdot a &= (x \cdot a) + (y \cdot a), \quad \forall x, y \in E, \forall a \in R, \\ x \cdot (ab) &= (x \cdot a) \cdot b, \quad \forall x \in E, \forall a, b \in R. \end{aligned}$$

If the ring R is unital, we will say that E is a **unital right R -module** if the additional property here below is satisfied:

$$x \cdot 1_R = x, \quad \forall x \in E.$$

Analogously, we can define a **(unital) left R -module** E over a ring R .

Note that a unital module over a unital complex algebra is naturally a complex vector space.

Definition 2.12. A **right pre-Hilbert C^* -module** M_B over a unital C^* -algebra B is a unital right module over the unital ring B that is equipped with a B -valued inner product $(x, y) \mapsto \langle x | y \rangle$ such that

$$\begin{aligned} \langle z | x + y \rangle_B &= \langle z | x \rangle_B + \langle z | y \rangle_B, \quad \forall x, y, z \in M, \\ \langle z | x \cdot b \rangle_B &= \langle z | x \rangle_B b, \quad \forall b \in B, \\ \langle y | x \rangle_B &= \langle x | y \rangle_B^*, \quad \forall x, y \in M, \\ \langle x | x \rangle_B &\in B_+, \quad \forall x \in M, \\ \langle x | x \rangle_B &= 0_B \Rightarrow x = 0_M. \end{aligned}$$

Analogously, a **left pre-Hilbert C^* -module** ${}_A M$ over a unital C^* -algebra A is a unital left module M over the unital ring A , that is equipped with an A -valued inner product $M \times M \rightarrow A$ denoted by $(x, y) \mapsto_A \langle x | y \rangle$. Here the A -linearity is on the first variable.

Remark 2.13. A right (respectively left) pre-Hilbert C^* -module M_B over the C^* -algebra B is naturally equipped with a norm

$$\|x\|_M := \sqrt{\|\langle x | x \rangle_B\|_B}, \quad \forall x \in M. \quad \lrcorner$$

Definition 2.14. A right (resp. left) **Hilbert C^* -module** is a right (resp. left) pre-Hilbert C^* -module over a C^* -algebra B that is a Banach space with respect to the previous norm $\|\cdot\|_M$ (resp. $\| \cdot \|_M$).

Definition 2.15. A right Hilbert C^* -module M_B is said to be **full** if

$$\overline{\langle M_B | M_B \rangle_B} := \overline{\text{span}\{\langle x | y \rangle_B \mid x, y \in M_B\}} = B,$$

where the closure is in the norm topology of the C^* -algebra B . A similar definition holds for a left Hilbert C^* -module.

We recall here the following well-known result, see for example in [GVF, Page 65]:

Lemma 2.16. Let M_B be a right Hilbert C^* -module over a unital C^* -algebra B . Then M_B is full if and only if $\text{span}\{\langle x | y \rangle_B \mid x, y \in M_B\} = B$.

We also recall a few definitions and results on bimodules (see for example [BCL2]).

Theorem 2.17. ([BCL2, Proposition 2.6]) Let M_A be a right unital Hilbert C^* -module over a unital C^* -algebra A and $J \subset A$ an involutive ideal in A .

The set $MJ := \{\sum_{j=1}^N x_j a_j \mid x_j \in M, a_j \in J, N \in \mathbb{N}_0\}$ is a submodule of M . The quotient module $M/(MJ)$ has a natural structure as a right Hilbert C^* -module over the quotient C^* -algebra A/J . If M is full over A , also $M/(MJ)$ is full over A/J .

A similar statement holds for a left Hilbert C^* -module.

Recall that a unital bimodule ${}_A M_B$ over two unital rings A and B is a left unital A -module and a right unital B -module such that $(a \cdot x) \cdot b = a \cdot (x \cdot b)$, for all $a \in A, b \in B$ and $x \in M$.

Definition 2.18. A **pre-Hilbert C^* -bimodule** ${}_A M_B$ over a pair of unital C^* -algebras A, B is a left pre-Hilbert C^* -module over A and a right pre-Hilbert C^* -module over B such that:

$$\begin{aligned} (a \cdot x) \cdot b &= a \cdot (x \cdot b), \quad \forall a \in A, x \in M, b \in B, \\ \langle x \mid ay \rangle_B &= \langle a^* x \mid y \rangle_B, \quad \forall x, y \in M, \forall a \in A, \\ {}_A \langle xb \mid y \rangle &= {}_A \langle x \mid yb^* \rangle, \quad \forall x, y \in M, \forall b \in B. \end{aligned}$$

A **Hilbert C^* -bimodule** ${}_A M_B$ is a pre-Hilbert C^* -bimodule over A and B that is simultaneously a left Hilbert C^* -module over A and a right Hilbert C^* -module over B . A full Hilbert C^* -bimodule over the C^* -algebras A_B is said to be an **imprimitivity bimodule** or an **equivalence bimodule** if:

$${}_A \langle x \mid y \rangle \cdot z = x \cdot \langle y \mid z \rangle_B \quad \forall x, y, z \in M.$$

Remark 2.19. (see for example [BCL2, Remark 2.14]) In an A - B pre-Hilbert C^* -bimodule there are two, usually different, norms, one as a left- C^* -module over A and one as a right- C^* -bimodule over B :

$${}_M \|x\| := \sqrt{\|{}_A \langle x \mid x \rangle\|_A}, \quad \|x\|_M := \sqrt{\|\langle x \mid x \rangle_B\|_B}, \quad \forall x \in M,$$

but this two norms coincide for an imprimitivity bimodule. In fact,

$$\begin{aligned} {}_M \|x\|^4 &= \|{}_A \langle x \mid x \rangle\|_A^2 = \|{}_A \langle x \mid x \rangle_A \langle x \mid x \rangle\|_A = \|{}_A \langle x \mid x \rangle_B \mid x \rangle\|_A \\ &\leq \|\langle x \mid x \rangle_B\|_B \cdot \|{}_A \langle x \mid x \rangle\|_A = \|x\|_M^2 \cdot \|x\|^2. \end{aligned} \quad \lrcorner$$

Definition 2.20. We say that a bimodule ${}_A M_A$ is a **symmetric bimodule** if $ax = xa$, for all $x \in M$ and all $a \in A$. If ${}_A M_A$ is a Hilbert C^* -bimodule, we say that it is a **symmetric C^* -bimodule** if it is symmetric as a bimodule and ${}_A \langle x \mid y \rangle = \langle y \mid x \rangle_A$, for all $x, y \in M$.

3 Main Results

First of all, we review the definition of Kreĭn C^* -algebra introduced by K.Kawamura [K2] and we explore some elementary properties of a Kreĭn C^* -algebra.

Definition 3.1. (K.Kawamura [K2, Definition 2]) A **Kreĭn C^* -algebra** is a Banach $*$ -algebra A admitting at least one **fundamental symmetry** i.e. a $*$ -automorphism $\phi : A \rightarrow A$ with $\phi \circ \phi = \iota_A$ such that $\|\phi(a^*)a\| = \|a\|^2$, for all $a \in A$.

Note that, in general, a Kreĭn C^* -algebra can admit several different fundamental symmetries.

Remark 3.2. A C^* -algebra A is a Kreĭn C^* -algebra with fundamental symmetry ι_A and so a Kreĭn C^* -algebra is a generalization of a C^* -algebra. Let (A, α) be a Kreĭn C^* -algebra with a given fundamental symmetry α . Then we always have the decomposition

$$A = A_+ \oplus A_-, \quad \text{where } A_+ = \{x \in A \mid \alpha(x) = x\}, \quad A_- = \{x \in A \mid \alpha(x) = -x\}.$$

Indeed, if $a \in A_+ \cap A_-$, then $a = \phi(a) = -a$ and so $a = 0$. Moreover for all $a \in A$, $a = \frac{a+\alpha(a)}{2} + \frac{a-\alpha(a)}{2}$ where $\frac{a+\alpha(a)}{2} \in A_+$ and $\frac{a-\alpha(a)}{2} \in A_-$. $\lrcorner$

Every Kreĭn C^* -algebra A with a given fundamental symmetry α becomes naturally a C^* -algebra, denoted here by $(A, \dagger_\alpha)$ when equipped with a new involution $\dagger_\alpha : A \rightarrow A$ as described in the following theorem.

Theorem 3.3. *Let (A, α) be a Kreĭn C^* -algebra with fundamental symmetry α . Then A becomes a C^* -algebra when equipped with the new involution $\dagger_\alpha$ defined by $\dagger_\alpha : x \mapsto \alpha(x)^*$. Furthermore, the fundamental symmetry α is an automorphism of this C^* -algebra and hence continuous in norm.*

Proof. It is straightforward from the definition of Kreĭn C^* -algebra to verify the first assertion. Since $\alpha(x^{\dagger_\alpha}) = \alpha(\alpha(x)^*) = \alpha(\alpha(x))^* = \alpha(x)^{\dagger_\alpha}$, for all $x \in A$, we have that α is a $*$ -homomorphism of the C^* -algebra $(A, \dagger_\alpha)$ into itself and hence it is continuous in norm. $\square$

Remark 3.4. Note that given a $\dagger$ -automorphism $\alpha : A \rightarrow A$ of a C^* -algebra A (with involution denoted by $\dagger$), we can naturally construct a Kreĭn C^* -algebra $(A, *_\alpha)$ with involution $x^{*_\alpha} := \alpha(x^\dagger)$, for $x \in A$ and that α becomes a fundamental symmetry for this Kreĭn C^* -algebra. $\lrcorner$

Now we observe the algebraic structure of the even and odd part of (A, α) .

Theorem 3.5. *Let (A, α) be a unital Kreĭn C^* -algebra with a given fundamental symmetry α . Then A_+ is a unital C^* -algebra and A_- is a unital Hilbert C^* -bimodule over A_+ .*

Proof. Note first that on A_+ the two involutions $*$ and $\dagger_\alpha$ coincide and since A_+ is closed under multiplication and involution, it is clearly a unital C^* -algebra.

We define right and left multiplications from $A_- \times A_+$ and $A_+ \times A_-$ into A_- as usual multiplications in A and we define a pair of A_+ -valued inner products from $A_- \times A_-$ into A_+ by

$$A_+ \langle x | y \rangle = xy^{\dagger_\alpha} \quad \text{and} \quad \langle x | y \rangle_{A_+} = x^{\dagger_\alpha} y \quad (3.1)$$

for all $x, y \in A_-$. With this definitions A_- has the structure of unital Hilbert C^* -bimodule on A_+ . $\square$

Additionally the Hilbert C^* -bimodule A_- has the property,

$$A_+ \langle x | y \rangle z = (xy^{\dagger_\alpha})z = x(y^{\dagger_\alpha} z) = x \langle y | z \rangle_{A_+}.$$

for all $x, y, z \in A_-$.

For any Kreĭn C^* -algebra equipped with a fundamental symmetry α , the odd part A_- in the fundamental decomposition $A = A_+ \oplus A_-$ is a Hilbert C^* -bimodule but it is not in general an imprimitivity bimodule because A_- might not be a full bimodule over A_+ . In the following we will usually assume this further property.

We will see later that the commutativity of A_+ and imprimitivity of A_- play important roles in order to get our spectral theory for Kreĭn C^* -algebras.

Definition 3.6. *Let (A, α) be a Kreĭn C^* -algebra with a given fundamental symmetry α . A Kreĭn C^* -algebra (A, α) is said to be **imprimitive** or **full** if its odd part A_- is an imprimitivity bimodule over A_+ . We say that (A, α) is **rank-one** if $\dim A_+ = \dim A_- = 1$ as complex vector spaces.*

Theorem 3.7. *Let (A, α) be a Kreĭn C^* -algebra with a given fundamental symmetry α . The algebra A is commutative if and only if the even part A_+ is a commutative unital C^* -algebra and the odd part is a symmetric Hilbert C^* -bimodule over A_+ . In particular a rank-one Kreĭn C^* -algebra is always commutative.*

Proof. If A is commutative, clearly A_- is a symmetric bimodule over the commutative algebra A_+ furthermore the inner products defined in (3.1) above satisfy $A_+ \langle x | y \rangle = xy^{\dagger_\alpha} = y^{\dagger_\alpha} x = \langle y | x \rangle_{A_+}$ and hence A_- is a symmetric C^* -bimodule over the commutative C^* -algebra A_+ . Conversely, if A_- is a symmetric C^* -bimodule, $a_- b_- = A_+ \langle a_- | b_-^{\dagger_\alpha} \rangle = \langle b_-^{\dagger_\alpha} | a_- \rangle_{A_+} = b_- a_-$, for all $a_-, b_- \in A_-$. Now, for all $a, b \in A$ with decompositions $a = a_+ + a_-$, $b = b_+ + b_-$ an easy computation shows: $ab = (a_+ + a_-)(b_+ + b_-) = a_+ b_+ + a_+ b_- + a_- b_+ + a_- b_- = b_+ a_+ + b_- a_+ + b_+ a_- + b_- a_- = ba$. $\square$

Definition 3.8. *We will say that a Kreĭn C^* -algebra is **symmetric** if there exists an **odd symmetry** i.e. a linear map $\epsilon : A \rightarrow A$ such that $\epsilon \circ \epsilon = i_A$, $\epsilon(x^*) = -\epsilon(x)^*$, for all $x \in A$, $\epsilon(xy) = \epsilon(x)y = x\epsilon(y)$, for all $x, y \in A$ and if there exists a fundamental symmetry α such that $\epsilon \circ \alpha = -\alpha \circ \epsilon$ (in this case we say that the symmetry and the odd symmetry are compatible).*

Lemma 3.9. *If ϵ is an odd symmetry of a symmetric Kreĭn C^* -algebra compatible with the symmetry α , then ϵ is always isometric.*

Proof. For all $x \in A$, $\|\epsilon(x)\|^2 = \|\epsilon(x)^{\dagger_\alpha} \epsilon(x)\| = \|\epsilon(x^{\dagger_\alpha}) \epsilon(x)\| = \|\epsilon \circ \epsilon(x^{\dagger_\alpha} x)\| = \|x^{\dagger_\alpha} x\| = \|x\|^2$. $\square$

Next we give an example of rank-one unital full symmetric Kreĭn C^* -algebra which has an important role in the proof of our spectral theory. Since, as we will see, in this case, the fundamental symmetry is necessarily unique, this is just a well-known example of commutative $\mathbb{Z}_2$ -graded C^* -algebra. Although all the properties described in the following three theorems are “standard” from the theory of $\mathbb{Z}_2$ -graded C^* -algebras, for the convenience of the reader we present here a direct proof of all of them in the “spirit” of Kreĭn C^* -algebras.

Theorem 3.10. *There is a rank-one unital symmetric Kreĭn C^* -algebra.*

Proof. For each $a, b \in \mathbb{C}$, let $T_{a,b} = \begin{bmatrix} a & b \\ b & a \end{bmatrix}$. Let $\mathbb{K} = \{T_{a,b} : a, b \in \mathbb{C}\}$ with the usual matrix operations of addition and multiplication. We define the involution by $T_{a,b}^* = T_{\bar{a}, \bar{b}}$. Furthermore, we equip $\mathbb{K}$ with the operator norm and choose as fundamental symmetry $\gamma : \mathbb{K} \rightarrow \mathbb{K}$ defined by $\gamma \begin{pmatrix} a & b \\ b & a \end{pmatrix} = \begin{pmatrix} a & -b \\ -b & a \end{pmatrix}$. Then we can write $\mathbb{K} = \mathbb{K}_+ \oplus \mathbb{K}_-$ where

$$\mathbb{K}_+ = \left\{ \begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix} : a \in \mathbb{C} \right\} \quad \text{and} \quad \mathbb{K}_- = \left\{ \begin{bmatrix} 0 & b \\ b & 0 \end{bmatrix} : b \in \mathbb{C} \right\}.$$

The symmetry of the algebra $\mathbb{K}$ can be checked defining $\epsilon(T_{a,b}) := T_{b,a}$. $\square$

Theorem 3.11. *The identity $\iota_{\mathbb{K}}$ and γ are the only two unital $*$ -automorphisms from $\mathbb{K}$ onto itself. Furthermore, γ is the unique fundamental symmetry on $\mathbb{K}$.*

Proof. Let ϕ be the unital $*$ -homomorphism from $\mathbb{K}$ into itself.

Suppose that $\phi(e) = \phi \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} a & b \\ b & a \end{pmatrix}$ for some $a, b \in \mathbb{C}$. Since $\phi(e)^2 = 1$ and $\phi(e^*) = -\phi(e)$, we have $(a, b) = (0, 1)$ or $(a, b) = (0, -1)$. Then $\phi = \iota_{\mathbb{K}}$ or γ , and this proves the theorem. $\square$

We can characterize the rank-one unital Kreĭn C^* -algebras as in Gel’fand-Mazur Theorem for C^* -algebras.

Theorem 3.12. *Every rank-one unital Kreĭn C^* -algebra is isomorphic to $\mathbb{K}$.*

Proof. By Gel’fand-Mazur Theorem, $A_+ = \mathbb{C}1$. Since $\dim A_- = 1$, there is a non-zero element e such that $\langle e \rangle = A_-$. WLOG, we choose e such that $\|e\| = 1$. Since $e^* \in A_-$ and $e \cdot e \in A_+$, we have $e^* = \alpha e$ and $e \cdot e = \beta$ for some $\alpha, \beta \in \mathbb{C}$. From properties in Kreĭn C^* -algebra, we can verify that $\alpha = \pm \exp(i\theta)$ and $\beta = \exp(i\theta)$ for some $\theta \in [0, 2\pi)$. Now we have that $e^* = \pm \exp(-i\theta)e$ and $e \cdot e = \exp(i\theta)$ for some $\theta \in [0, 2\pi)$. We use the notation $A_{\pm\theta}$ corresponding with the preceding operations. Since $\|x\| = \|L_x\|$ where L_x is the left multiplication operator, we can prove that $\|x\| = \max\{|a+b|, |a-b|\}$. It is easy to show that θ must be zero or π .

For case $\theta = \pi$, we let $x = y = i + e$, then

$$\|xy\| = \max\{|-2+2i|, |-2-2i|\} = 2\sqrt{2} \not\leq 2 = \max\{|i+1|^2, |i-1|^2\} = \|x\|\|y\|.$$

Hence $A_{\pm\pi}$ is not a Kreĭn C^* -algebra.

For A_{-0} , we have $e \cdot e = 1$ and $e^* = -e$. Since every element in A_{0-} is in the form $m1 + ne$, where m and n are in $\mathbb{C}$, we define $f : A_{0-} \rightarrow \mathbb{K}$ by $f(1) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $f(e) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$. Then f is an isomorphism between the Kreĭn C^* -algebras A_{0-} and $\mathbb{K}$. Since $\mathbb{K}$ is Kreĭn C^* -algebra, A_{-0} is a Kreĭn C^* -algebra too.

For $A_{+\theta}$, we have $e \cdot e = 1$ and $e^* = e$. Suppose that γ is the fundamental symmetry on it. Note that $\gamma(x^{*-}) = x^{*+}$ for all $x \in A$ where $*_-$ and $*_+$ are involutive on $A_{-\theta}$ and $A_{+\theta}$, respectively. Then

$$\|x\|^2 = \|\gamma(x^{*+})x\| = \|x^{*-}x\|,$$

which contradicts to the fact that A_- is the nontrivial Kreĭn C^* -algebra. Since γ is the only possible fundamental symmetry on A , we can conclude that A_+ is not a Kreĭn C^* -algebra. $\square$

Theorem 3.13. *The space $C(M, K)$ of all continuous functions from a compact Hausdorff space M into a Kreĭn C^* -algebra K is a unital Kreĭn C^* -algebra.*

Proof. Let γ be a fundamental symmetry of the Kreĭn C^* -algebra K and M a compact Hausdorff space. We define all the operations and the norm as follows:

$$\begin{aligned} (f+g)(x) &= f(x) + g(x) & (fg)(x) &= f(x)g(x) \\ (kf)(x) &= kf(x) & f^{*C}(x) &= f(x)^{*K} \\ \|f\|_C &= \sup_{x \in M} \|f(x)\|_K, \end{aligned}$$

for all $k \in \mathbb{C}$, $x \in M$ and for all $f, g \in C(M, K)$. It is easy to check that $C(M, K)$ is a Banach algebra and a $*$ -algebra with the operations and norm above.

To see that $C(M, K)$ is a Kreĭn C^* -algebra, consider the map ϕ_C from $C(M, K)$ into itself defined by $\phi_C(f) = \gamma \circ f$. Since, for all $x \in M$,

$$\begin{aligned} \phi_C(fg)(x) &= \gamma \circ (fg)(x) = \gamma(f(x)g(x)) = \gamma(f(x))\gamma(g(x)) = \phi_C(f)\phi_C(g)(x) \\ \phi_C(f^{*C})(x) &= (\gamma \circ f^{*C})(x) = \gamma(f(x)^{*K}) = \gamma(f(x))^{*K} \\ &= (\gamma \circ f)(x)^{*K} = (\gamma \circ f)^{*C}(x) = \phi_C(f)^{*C}(x), \\ \phi_C(1_C)(x) &= \gamma(1_C(x)) = \gamma(1_K) = 1_K = 1_C(x), \end{aligned}$$

ϕ_C is a unital $*$ -homomorphism that is an involutive $*$ -automorphism because

$$\phi_C \circ \phi_C(f) = \phi_C(\gamma \circ f) = \gamma \circ \gamma \circ f = f, \quad \forall f \in C(M, K),$$

and also a fundamental symmetry of a Kreĭn C^* -algebra because

$$\begin{aligned} \|\phi_C(f)^{*C} f\|_C &= \sup_{x \in M} \|(\gamma \circ f)^{*C}(x) f(x)\|_K = \sup_{x \in M} \|(\gamma \circ f)(x)^{*K} f(x)\|_K \\ &= \sup_{x \in M} \|\gamma(f(x))^{*K} f(x)\|_K = \sup_{x \in M} \|f(x)\|_K^2 = \|f\|_C^2, \end{aligned}$$

for all $f \in C(M, K)$. $\square$

When we take $K := \mathbb{K}$ in the previous theorem, we obtain another example of commutative symmetric imprimitivity Kreĭn C^* -algebra.

Corollary 3.14. *The Kreĭn C^* -algebra $C(M, \mathbb{K})$ is a commutative symmetric imprimitivity unital Kreĭn C^* -algebra.*

Proof. Define the fundamental symmetry $\phi_C(f) := \gamma \circ f$ and $\epsilon_C(f) := \epsilon_{\mathbb{K}} \circ f$. $\square$

Theorem 3.15. *Let (A, α) be a (unital commutative) Kreĭn C^* -algebra with a fundamental symmetry α . Let I be a closed ideal in A invariant under α , i.e. $\alpha(I) \subseteq I$, then A/I is also a (unital commutative) Kreĭn C^* -algebra.*

Proof. Since I is a closed ideal in the C^* -algebra $(A, \dagger_\alpha)$, the quotient $(A/I, \dagger_\alpha)$ is a C^* -algebra with involution $(x+I)^{\dagger_\alpha} := x^{\dagger_\alpha} + I$. Since I is invariant under the $*$ -automorphism α we can define $[\alpha] : A/I \rightarrow A/I$ by $x+I \mapsto \alpha(x)+I$ for all $x \in A$ and $[\alpha]$ is a $\dagger_\alpha$ -automorphism of $(A/I, \dagger_\alpha)$. By remark 3.4 $(A/I, *)$ is a Kreĭn C^* -algebra with involution $(x+I)^* := \alpha(x^{\dagger_\alpha}) + I = x^* + I$ and $[\alpha]$ becomes the fundamental symmetry on $(A/I, *)$. $\square$

Theorem 3.16. Let (A, α) and (B, β) be two Kreĭn C^* -algebras with given fundamental symmetries α and β . A $*$ -homomorphism $\phi : A \rightarrow B$ satisfying the property $\phi \circ \alpha = \beta \circ \phi$ is always continuous. Additionally, $\phi(A_+) \subseteq B_+$ and $\phi(A_-) \subseteq B_-$.

Proof. Note that ϕ is a $\dagger$ -homomorphism between the associated C^* -algebras $(A, \dagger_\alpha)$ and $(B, \dagger_\beta)$. The “invariance” of ϕ under α, β implies the last property. $\square$

If $\phi : A \rightarrow B$ is a unital $*$ -homomorphism such that $\phi \circ \alpha = \beta \circ \phi$ for a given pair of fundamental symmetries of the Kreĭn C^* -algebras A and B , in view of the last property in theorem 3.16, we will denote by $\phi_+ : A_+ \rightarrow B_+$ and $\phi_- : A_- \rightarrow B_-$ the restrictions to the even and odd parts of the Kreĭn C^* -algebras. Note that $\phi = \phi_+ \oplus \phi_-$. In particular, the quotient isomorphism $\pi : A \rightarrow A/I$ from a Kreĭn C^* -algebra A to its quotient Kreĭn C^* -algebra by an ideal I that is invariant under a fundamental symmetry α of A , can be written as a direct sum $\pi = \pi_+ \oplus \pi_-$ of the isomorphisms $\pi_+ : A_+ \rightarrow (A/I)_+$ and $\pi_- : A_- \rightarrow (A/I)_-$.

Corollary 3.17. Let (A, α) be a unital Kreĭn C^* -algebra with a fundamental symmetry α . Let w be a unital $*$ -homomorphism from A into $\mathbb{K}$ with the property that $w \circ \alpha = \gamma \circ w$. Then $A/\ker(w)$ is a Kreĭn C^* -algebra.

Proof. By the preceding theorem, $\ker(w)$ is a closed ideal in A . A direct computation shows that $\alpha(\ker(w)) \subseteq \ker(w)$. Consequently, $A/\ker(w)$ is a Kreĭn C^* -algebra by theorem 3.15. $\square$

Corollary 3.18. Let (A, α) be a unital imprimitive Kreĭn C^* -algebra with a fundamental symmetry α . Let w be a unital $*$ -homomorphism from A into $\mathbb{K}$ with the property that $w \circ \alpha = \gamma \circ w$. Then $A/\ker(w) \cong \mathbb{K}$.

Proof. $A/\ker(w)$ is a Kreĭn C^* -algebra by Corollary 3.17.

Define $\beta : A/\ker(w) \rightarrow \mathbb{K}$ by $x + \ker(w) \mapsto w(x)$ for all $x \in A$. It is easily checked that β is unital injective $*$ -homomorphism. It remains to show the surjective property. Since A_{-A_+} is full, WLOG $1 = \sum_{j=1}^n x_j y_j$ for some $n \in \mathbb{N}$ and $x_j, y_j \in A_-$ for all $j = 1, \dots, n$. Suppose that $w(x) = 0$ for all $x \in A_-$. Since w is a unital homomorphism, we have

$$1 = w(1) = w\left(\sum_{j=1}^n x_j y_j\right) = \sum_{j=1}^n w(x_j)w(y_j) = 0,$$

which leads to contradiction. Thus there is element x in A_- such that $w(x) \neq 0$. Since $w(A_-) \subseteq \mathbb{K}_-$ and $\dim \mathbb{K}_- = 1$, we get $w(A_-) = \mathbb{K}_-$. Similarly, by Gel’fand-Mazur theorem $w(A_+) = \mathbb{K}_+$. Thus we already have

$$\beta(A/\ker(w)) = w(A) = w(A_+ \oplus A_-) = w(A_+) \oplus w(A_-) = \mathbb{K}_+ \oplus \mathbb{K}_- = \mathbb{K},$$

and so β is clearly surjective. $\square$

Definition 3.19. A **character** on a unital (commutative symmetric full) Kreĭn C^* -algebra A is a unital $*$ -homomorphism $w : A \rightarrow \mathbb{K}$ such that there exists at least one fundamental symmetry α of the Kreĭn C^* -algebra A satisfying the property $w \circ \alpha = \gamma \circ w$. In this case we will say that α and w are **compatible**.

We denote by $\Omega(A)$ the set of characters on A , and by $\Omega(A, \alpha)$ the set of characters compatible with a given fundamental symmetry α that is

$$\Omega(A, \alpha) = \{w \mid w : A \rightarrow \mathbb{K} \text{ unital } * \text{-homomorphism compatible with } \alpha\}.$$

For each $a \in A$, define $\hat{a} : \Omega(A) \rightarrow \mathbb{K}$ by $\hat{a}(w) = w(a)$ for all $w \in \Omega(A)$. Equip $\Omega(A)$ with the smallest topology which makes each $\hat{a}$ continuous.

Note that $\Omega(A, \alpha)$ with the induced subspace topology is clearly a closed set in the topological space $\Omega(A)$ since $\Omega(A, \alpha) = \{w \in \Omega(A) \mid \alpha(\hat{x})(w) = \gamma \circ \hat{x}(w), \forall x \in A\}$ and the functions $\alpha(\hat{x}), \gamma \circ \hat{x}$ are continuous on $\Omega(A)$.

We next define an equivalence relation between the characters as follows:

$$w_1 \sim w_2 \iff w_2 = w_1 \circ \phi \text{ for some } \phi \text{ a unital } *- \text{automorphism on } \mathbb{K}.$$

By Theorem 3.11, $[w] = \{w, \gamma \circ w\}$. Note that $w \in \Omega(A, \alpha)$ implies $[w] \subset \Omega(A, \alpha)$.

$$\text{We define } \Omega_b(A) = \{[w] \mid w \in \Omega(A)\}, \quad \Omega_b(A, \alpha) = \{[w] \mid w \in \Omega(A, \alpha)\}.$$

Equip $\Omega_b(A)$ with the quotient topology induced by the quotient map

$$\mu : \Omega(A) \rightarrow \Omega_b(A) \text{ given by } \mu : w \mapsto [w].$$

Note that on $\Omega_b(A, \alpha)$ the quotient topology induced by $\Omega(A, \alpha)$ coincides with the subspace topology induced by $\Omega_b(A)$.

Lemma 3.20. $\Omega(A)$ and $\Omega(A, \alpha)$ are compact Hausdorff spaces.

Proof. Since a character $w : A \rightarrow \mathbb{K}$ is a unital $*$ -homomorphism of the associated C^* -algebras $(A, \dagger_\alpha)$ and $(\mathbb{K}, \dagger_\gamma)$, with the same techniques used in Banach-Alaoglu theorem, it is a standard matter to check that $\Omega(A)$ is a closed subset of compact set $\prod_{x \in A} \overline{B(0_{\mathbb{K}}, \|x\|)}$ and hence $\Omega(A)$ is also a compact set. To show that $\Omega(A)$ is a Hausdorff space, we first let w_1, w_2 be characters such that $w_1 \neq w_2$. Then there is $a \in A$ such that $w_1(a) \neq w_2(a)$ that is $\hat{a}(w_1) \neq \hat{a}(w_2)$. Since $\hat{a}$ is continuous on $\Omega(A)$ and $\mathbb{K}$ is Hausdorff, we obtain that $\Omega(A)$ is also Hausdorff. Since $\Omega(A, \alpha)$ is a closed subspace of $\Omega(A)$, the result follows. $\square$

Lemma 3.21. Let w, w_1 and w_2 be characters. The following properties hold.

- a) If $w_1 \sim w_2$, then $\ker(w_1) = \ker(w_2)$.
- b) If $w \in \Omega(A, \alpha)$, $w(x) = 0 \iff w(x^{\dagger_\alpha} x) = 0$, for all $x \in A$.
- c) If $w_1, w_2 \in \Omega(A, \alpha)$ and $w_{1+} = w_{2+}$, then $\ker(w_1) = \ker(w_2)$.

Proof. Property a) is clear. For b), assume that $w(x^{\dagger_\alpha} x) = 0$. Then

$$\|w(x)\|^2 = \|\gamma(w(x))^* w(x)\| = \|w(\alpha(x))^* w(x)\| = \|w(\alpha(x^*)x)\| = \|w(x^{\dagger_\alpha} x)\| = 0,$$

and so $w(x) = 0$.

For c), by the assumption, we have $\ker(w_{1+}) = \ker(w_{2+})$. Let $x \in A_-$ be such that $w_1(x) = 0$. By b), $w_1(\alpha(x^*)x) = 0$ and also $w_2(\alpha(x^*)x) = 0$ because $\ker(w_{1+}) = \ker(w_{2+})$. Again by b), $w_2(x) = 0$ and hence $\ker(w_{1-}) \subseteq \ker(w_{2-})$. By the same argument, it is elementary to verify the inverse inclusion. $\square$

Lemma 3.22. Let (A, α) be a Kreĭn C^* -algebra equipped with a fundamental symmetry α and let $\omega \in \Omega(A_+)$ be a character defined on the even part of the Kreĭn C^* -algebra $A = A_+ \oplus A_-$. Define

$$I_+ := \ker(\omega), \quad I_- := A_- \ker(\omega) := \text{span}\{xa \mid x \in A_-, a \in \ker \omega\}, \quad I := I_+ \oplus I_-.$$

Then I is an ideal in A invariant under α and the following properties hold

- a) A_+/I_+ is a C^* -algebra with dimension 1.
- b) A_-/I_- is a Hilbert C^* -bimodule over A_+/I_+ .
- c) $A/I = (A/I)_+ \oplus (A/I)_- \cong A_+/I_+ \oplus A_-/I_-$.

d) A/I is a rank-one Kreĭn C^* -algebra.

Proof. The bimodule A_-/I_- over A_+/I_+ is a Hilbert C^* -bimodule with the inner products defined as in the proof of theorem 3.5 using the $(x + I_-)^\dagger := x^\dagger + I_-$ involution. The only other thing that is not completely straightforward is that A/I is rank 1. By b), since by Gel'fand-Mazur $A_+/I_+ \cong \mathbb{C}$, A_-/I_- is a Hilbert space over A_+/I_+ . To show the rank-one property, suppose by contradiction that $x, y \in A_-/I_-$ is a pair of orthonormal vectors, then

$$y = \langle x|x \rangle y + \langle y|x \rangle y = \langle x+y|x \rangle y = (x+y)\langle x|y \rangle = 0,$$

which is impossible. $\square$

Theorem 3.23. *If (A, α) is a unital commutative imprimitive Kreĭn C^* -algebra with fundamental symmetry α , then $\Omega_b(A, \alpha)$ is a compact Hausdorff space.*

Proof. Since $\Omega(A, \alpha)$ is compact by lemma 3.20 and μ is continuous, $\Omega_b(A, \alpha)$ is also compact. We now consider the map ϕ from $\Omega_b(A, \alpha)$ to $\Omega(A_+)$ defined by $[w] \mapsto w_+$. If we can show that this map is a homeomorphism, we can conclude that $\Omega_b(A, \alpha)$ is a Hausdorff space since $\Omega(A_+)$ is a Hausdorff space by the spectral theorem for unital commutative C^* -algebras. It is sufficient to show that ϕ is a continuous bijective map because $\Omega_b(A, \alpha)$ is a compact space and $\Omega(A)$ is a Hausdorff space.

Firstly, ϕ is well-defined. More precisely, let $[w_1] = [w_2]$, if $w_1 \neq w_2$, then $w_2 = \gamma \circ w_1$. Let $x \in A_+$, then $w_1(x) = w_1(\alpha(x)) = \gamma(w_1(x)) = w_2(x)$, that is, $w_{1+} = w_{2+}$.

We next show that $R := \phi \circ \mu$ is a continuous map. Note that, by definition, we have $R(w) = w_+$. For easier consideration, we provide a diagram of all the functions involved,

$$\begin{array}{ccccc} & & \Omega(A, \alpha) & & \\ & \swarrow \mu & \downarrow R & \searrow \hat{a} & \\ \Omega_b(A, \alpha) & \xrightarrow{\phi} & \Omega(A_+) & \xrightarrow{\hat{a}_+} & \mathbb{C} \end{array}$$

since for all $a \in A_+$, $\hat{a} = \hat{a}_+ \circ R$ is continuous on $\Omega(A, \alpha)$, R is also continuous. Since R is a continuous map, by the properties of quotient topology we also have that ϕ is a continuous map. Next, we will show that ϕ is a bijection.

To show that ϕ is injective, we suppose that w_1, w_2 are characters on (A, α) such that $w_{1+} = w_{2+}$. We next examine the diagram here

$$\begin{array}{ccccc} A & \xrightarrow{w_1} & \mathbb{K} & \xleftarrow{w_2} & A \\ \pi_1 \downarrow & & & & \downarrow \pi_2 \\ A/\ker(w_1) \cong \mathbb{K} & \xrightarrow{\beta_1} & & \xleftarrow{\beta_2} & A/\ker(w_2) \cong \mathbb{K} \end{array}$$

By lemma 3.21 (c) and corollary 3.18 we have $A/\ker(w_1) \cong \mathbb{K} \cong A/\ker(w_2)$, $\beta_2 \circ \beta_1^{-1}$ is a unital $*$ -automorphism on $\mathbb{K}$. Since by theorem 3.11 a unital $*$ -automorphism on $\mathbb{K}$ is either the identity map or γ , we have to consider two cases.

1. $\beta_2 \circ \beta_1^{-1}$ is an identity on $\mathbb{K}$

Then $\beta_1 = \beta_2$. Since $\ker(w_1) = \ker(w_2)$, we have

$$w_1(a) = \beta_1(a + \ker w_1) = \beta_2(a + \ker w_2) = w_2(a),$$

for all $a \in A$. Thus $w_1 = w_2$.

$$2. \beta_2 \circ \beta_1^{-1} = \gamma$$

Then $\beta_2 = \gamma \circ \beta_1$. Again by the fact that $\ker(w_1) = \ker(w_2)$, we have

$$w_2(a) = \beta_2(a + \ker w_2) = \gamma \circ \beta_1(a + \ker w_1) = \gamma \circ w_1(a),$$

for all $a \in A$. Hence $w_2 = \gamma \circ w_1$.

From both of two cases, we can conclude that $[w_1] = [w_2]$ which implies the injection of ϕ , as we wanted. To prove the surjectivity of ϕ , let $w^\circ : A_+ \rightarrow \mathbb{K}_+$. Consider $I_+ := \ker(w^\circ)$ and define the ideal I in A as in lemma 3.22.

Since A/I is a rank-one Kreĭn C^* -algebra, A/I is isomorphic to $\mathbb{K}$ by an isomorphism f such that $f \circ [\alpha] = \gamma \circ f$, where $[\alpha]$ is the fundamental symmetry of A/I such that $\pi \circ \alpha = [\alpha] \circ \pi$. Note that $f = f_+ \oplus f_-$ where $f_+ : (A/I)_+ \rightarrow \mathbb{K}_+$ is the isomorphism $f_+(a + I) = w^\circ(a)$, for all $a \in A_+$.

$$A \xrightarrow{\pi} A/I \xrightarrow{f} \mathbb{K}.$$

Define $w := f \circ \pi$. Then $w \in \Omega(A, \alpha)$ because $w \circ \alpha = \gamma \circ w$. We claim that $w_+ = w^\circ$. To see this, let $a \in A_+$. Then $w(a) = f \circ \pi(a) = f(a + I) = w^\circ(a)$, and the theorem is proved. $\square$

Definition 3.24. If (A, α, ϵ) is a commutative full symmetric unital Kreĭn C^* -algebra with a given fundamental symmetry α and a given odd symmetry ϵ , we define the **even spectrum** as the set of **even characters** of A : $\Omega(A, \alpha, \epsilon) := \{w \in \Omega(A, \alpha) \mid \epsilon_{\mathbb{K}} \circ w \circ \epsilon = w\}$.

Theorem 3.25. If (A, α, ϵ) is a commutative unital full symmetric Kreĭn C^* -algebra with fundamental symmetry α and odd symmetry ϵ , the character $w \in \Omega(A, \alpha)$ is even if and only if $\gamma \circ w \in \Omega(A, \alpha)$ is not even. Hence in every equivalence class $[w] = \{w, \gamma \circ w\}$ there is one and only one even character and there is a bijection between $\Omega(A, \alpha, \epsilon)$ and $\Omega_b(A, \alpha)$.

Proof. Since $\epsilon_{\mathbb{K}} \circ \gamma = -\gamma \circ \epsilon_{\mathbb{K}}$, we see that w is even if and only if $\gamma \circ w$ is odd. Let w° be a character on A_+ and define $w(x) := w^\circ(x_+) + \epsilon_{\mathbb{K}} \circ w^\circ \circ \epsilon(x_-)$. By the properties of the odd symmetry, we see that w is an even character compatible with α . $\square$

Definition 3.26. Let A be a commutative unital full symmetric Kreĭn C^* -algebra and let α be a fundamental symmetry of A and ϵ an odd symmetry of A . The **Gel'fand transform** of $x \in A$ is the map $\hat{x} : \Omega_b(A, \alpha) \rightarrow \mathbb{K}$ defined by:

$$\hat{x}([w]) := w(x_+) + \epsilon_{\mathbb{K}} \circ w \circ \epsilon(x_-), \quad \forall x \in A.$$

By the previous theorem, it is clear that the Gel'fand transform $\hat{x}$ of x is just the function that to every even character $w \in \Omega(A, \alpha, \epsilon)$ associates $w(x)$.

Although the following theorem is our goal, the proof is easy and straightforward.

Theorem 3.27 (Spectral theorem). If (A, α, ϵ) is a unital commutative symmetric imprimitive Kreĭn C^* -algebra with fundamental symmetry α and odd symmetry ϵ , then the Gelfand transform

$$\varphi : A \rightarrow C(\Omega_b(A, \alpha), \mathbb{K}), \quad a \mapsto \hat{a}$$

is an isometric $*$ -isomorphism.

Proof. In view of theorem 3.25 let us denote by $[w]$ with w even a point of $\Omega_b(A, \alpha)$. To prove that φ is a $*$ -homomorphism of algebras, let $a, b \in A$ and $k \in \mathbb{C}$,

$$\begin{aligned} (\varphi(ab))([w]) &= (\widehat{ab})([w]) = w(ab) = w(a)w(b) = \hat{a}([w])\hat{b}([w]) = (\varphi(a)\varphi(b))([w]), \\ (\varphi(ka + b))([w]) &= (\widehat{ka + b})([w]) = w(ka + b) = kw(a) + w(b) = k\hat{a}([w]) + \hat{b}([w]) \\ &= (k\varphi(a) + \varphi(b))([w]), \\ \varphi(a^*)[w] &= \widehat{a^*}([w]) = [w](a^*) = w(a)^* = \varphi(a)^*[w]. \end{aligned}$$

for all $[w] \in \Omega_b(A, \alpha)$. Clearly $\varphi(1_A) = 1_C$ so that φ is unital.

It is easy to verify that $\varphi \circ \alpha = \phi_C \circ \varphi$. In fact, for all $a \in A$, $[w] \in \Omega_b(A, \alpha)$,

$$\begin{aligned} (\varphi \circ \alpha)(a)[w] &= \varphi(\alpha(a))[w] = w(\alpha(a)) = \gamma(w(a)) = \gamma(\varphi(a)[w]) \\ &= (\gamma \circ \varphi(a))[w] = \phi_C(\varphi(a))[w] = (\phi_C \circ \varphi)(a)[w]. \end{aligned}$$

We also have that $\varphi \circ \epsilon = \epsilon_C \circ \varphi$, in fact, for all $[w] \in \Omega_b(A, \alpha)$, and all $x \in A$,

$$\epsilon_C \circ \varphi(x)[w] = \epsilon_{\mathbb{K}}(w(x)) = \epsilon_{\mathbb{K}} \circ w \circ \epsilon^2(x) = w \circ \epsilon(x) = \varphi \circ \epsilon(x)[w]. \quad (3.2)$$

Let A be a unital commutative Kreĭn C^* -algebra with the fundamental symmetry α . Then $(A, \dagger_A)$ and $(C(\Omega_b(A, \alpha), \mathbb{K}), \dagger_C)$ become C^* -algebras with the involutions $\dagger_A$ and $\dagger_C$ defined as in theorem 3.3 respectively, that is,

$$a^{\dagger_A} = \alpha(a^{*A}) \text{ for all } a \in A \text{ and } f^{\dagger_C} = \phi_C(f^{*C}) \text{ for all } f \in C(\Omega_b(A, \alpha), \mathbb{K}).$$

To show φ is a $\dagger$ -homomorphism, let $a \in A$ and $[w] \in \Omega_b(A, \alpha)$,

$$\varphi(a^{\dagger_A})[w] = \varphi(\alpha(a^{*A}))[w] = \widehat{\alpha(a^{*A})}[w] = w(\alpha(a^{*A})).$$

Since $w \circ \alpha = \gamma \circ w$, we have,

$$\varphi(a^{\dagger_A})[w] = \gamma(w(a^{*A})) = \gamma(w(a)^{*_{\mathbb{K}}}) = \gamma(\hat{a}([w])^{*_{\mathbb{K}}}) = \gamma(\hat{a}^{*C}([w]))$$

Since $\phi_C(f) = \gamma \circ f$ for all $f \in C(\Omega_b(A, \mathbb{K}))$,

$$\varphi(a^{\dagger_A})[w] = \phi_C(\hat{a}^{*C})[w] = \hat{a}^{\dagger_C}[w] = \varphi(a)^{\dagger_C}[w].$$

By the spectral theorem for unital commutative C^* -algebras, the restriction of the Gel'fand transform to the even part $\varphi_+ : A_+ \rightarrow C(\Omega(A, \alpha), \mathbb{K})_+$ is an isometric $\dagger$ -isomorphism, for all $a \in A$ that coincides with the usual Gel'fand isomorphism for the commutative unital C^* -algebra A_+ .

Since ϵ and ϵ_C are linear surjective (because $\epsilon^2 = i_A$) isometries, from the equation (3.2), we see that $\varphi_- = \epsilon_- \circ \varphi_+ \circ \epsilon_{C+}$ is isometric surjective too and hence $\varphi = \varphi_+ \oplus \varphi_-$ is a surjective isometry that is also a $*$ -homomorphism. $\square$

The previous theorem provide a complete characterization of those unital commutative Kreĭn C^* -algebras that are full and symmetric. In this case, a posteriori, with a bit more work, we might actually prove that the fundamental symmetry and the odd symmetry are indeed unique. Further analysis is required in order to provide a spectral theory of more general Kreĭn C^* -algebras. Omitting the exchange symmetry requirement will lead us to algebras of sections of bundles of one-dimensional Kreĭn C^* -algebras and omitting the commutativity (or just the imprimitivity condition on the odd part) will lead us to a theory of Kreĭn spaceoids along very similar lines to those used to describe the spectrum of commutative full C^* -categories in [BCL1], we hope that the extra effort paid here to describe a “direct proof” of the spectral theory in the special case of commutative full symmetric Kreĭn C^* -algebras will facilitate the analysis of those more general topics that we plan to address in the near future.

Acknowledgements : We thank Dr. Roberto Conti for making several suggestions and providing references to arguments in the theory of C^* -dynamical systems ($\mathbb{Z}_2$ -graded C^* -algebras) that might be used to shortcut/simplify most of the proofs and results provided in this work. We plan to return on this topic and to improve the manuscript in this direction.

References

- [BCL1] Bertozzini P, Conti R, Lewkeeratiyutkul W (2008) A Horizontal Categorification of Gel'fand Duality [arXiv:0812.3601v1](#)¹
- [BCL2] Bertozzini P, Conti R, Lewkeeratiyutkul W (2008) A Spectral Theorem for Imprimitivity C^* -bi-modules [arXiv:0812.3596v1](#)
- [B] Bleuler K (1950) Eine Neue Methode zur Behandlung der Longitudinalen und Skalaren Photonen *Helv Phys Acta* 23(5):567-586
- [C] Connes A (1994) *Noncommutative Geometry* Academic Press
- [D] Dirac P (1942) The Physical Interpretation of Quantum Mechanics *Pro Roy Soc London Ser A* 180:1-40
- [DR] Ditschel M, Rovnyak J (1996) Operators on Indefinite Inner Product Spaces *Lectures on Operator Theory and its Application* Fields Institute Monographs 3:141-232 American Mathematical Society
- [G] Ginzburg J (1957) On J -contractive Operator Functions *Dokl Akad Nauk SSSR* 117:171-173
- [GVF] Gracia-Bondía J-M, Várilly J, Figueroa H (2001) *Elements of Non-commutative Geometry* Birkhäuser
- [Gu] Gupta S (1950) Theory of Longitudinal Photons in Quantum Electrodynamics *Proc Phys Soc A* 63(27):681-691
- [K1] Kawamura K (2006) Indefinite-metric Quantum Field Theory and Operator Algebra [arXiv:math/0608076v2](#)
- [K2] Kawamura K (2006) Algebra with Indefinite Involution and Its Representation in Kreĭn Space [arXiv:math/0610059v2](#)
- [M] Minkowski H (1909) Raum und Zeit *Physikalische Zeitschrift* 10:104-111
- [Mu] Murphy G (1990) *C^* -algebras and Operator Theory* Academic Press
- [PRV] Paschke M, Rennie A, Verch R, Lorentzian Spectral Triples, in preparation²
- [P] Pauli W (1943) On Dirac's New Method of Field Quantization *Rev Modern Phys* 15:175-207
- [Po] Pontrjagin L (1944) Hermitian Operators in Spaces with Indefinite Metric *Izv Akad Nauk SSSR Ser Mat* 8:243-280
- [S] Scheibe E (1960) Über Hermitesche Formen in Topologischen Vektorräumen I, *Ann Acad Fenn Ser A I* 294 30 pages
- [St] Strohmaier A (2006) On Noncommutative and Pseudo-Riemannian Geometry *J Geom Phys* 56(2):175-195 [arXiv:math-ph/0110001](#)

¹Now published as: Bertozzini P, Conti R, Lewkeeratiyutkul W (2011) A Horizontal Categorification of Gel'fand Duality *Adv Math* 226(1):584-607 [arXiv:0812.3601v2](#).

²Now published as: van den Dungen K, Paschke M, Rennie A (2013) Pseudo-Riemannian Spectral Triples and the Harmonic Oscillator *J Geom Phys* 73:37-55 [arXiv:1207.2112](#).